

Preface to special section on Global Aerosol System

D. Tanré,¹ Y. Kaufman,² T. Nakajima,³ and V. Ramanathan⁴

Published 26 March 2005.

Citation: Tanré, D., Y. Kaufman, T. Nakajima, and V. Ramanathan (2005), Preface to special section on Global Aerosol System, *J. Geophys. Res.*, 110, D10S01, doi:10.1029/2004JD005724.

[1] This special issue brings together papers that were presented in a workshop of experts on the global aerosol system and its impacts on climate. Aerosols originate as urban haze and rural haze, and owing to long range transport of pollutants, becomes widespread aerosol layers across the globe.

[2] Atmospheric aerosols play a dual role in the climate system and the hydrologic cycle (see reviews in the work of *Ramanathan et al.* [2001] and *Kaufman et al.* [2002]). Their first role is interaction with solar and thermal radiation: by scattering sunlight and reflecting a fraction of it back to space aerosols cool the climate system; by absorbing sunlight in the atmosphere aerosols further cool the surface but warm the atmosphere, changing the temperature and humidity profiles and the conditions for cloud development.

[3] Aerosol particles play a second role by affecting the hydrologic cycle: serving as cloud condensation nuclei and ice nuclei, aerosols control the cloud droplet concentration and size. Higher number concentrations of submicron pollution aerosol tend to decrease the cloud droplet size and prevent or delay development of precipitation. Supermicron sea-salt particles can reintroduce precipitation and oppose the effect of pollution aerosol. These processes may cause changes in precipitation patterns to which the human civilization adapted during the last millennium, changes in cloud cover, and possible changes in the frequency of extreme events.

[4] No less important is the aerosol effect on health, air quality, and agriculture that brought policy responses in Europe and North America in the last century and is affecting policy decisions in east and southern Asia today. Therefore advancements in aerosol science, together with research on the effect of greenhouse and trace gases and increased ability to predict the impact of human activity on the regional and global aerosol, can be expected to have direct impact on energy use and economic activities in nations around the world.

[5] Unraveling these aerosol effects is still difficult because of the high spatial and temporal variability of sources,

complex interaction of aerosol with atmospheric trace gases and clouds, and the short lifetime of aerosol of about a week, owing to wet and dry depositions. A single aerosol particle over the Mediterranean Sea can include anthropogenic sulfate and carbonaceous material from Europe, coating natural dust particle from Africa.

[6] Today's aerosol assessments of climate are based on aerosol chemical transport models and climate models that cannot yet resolve the aerosol interaction with clouds with their microscale physical and chemical exchanges. Models make good progress in capturing aerosol evolution, but source characterization is difficult. Accurate assessments of aerosol distribution and composition, the anthropogenic component, and the impacts on radiation and water cycle therefore require orchestrated observations of aerosol, precursor trace gases, clouds and precipitation from satellites, networks of surface-based instruments, laboratory studies, and dedicated field experiments. The observations require sufficient information content, accuracy, and global coverage to permit their integration into and testing of aerosol chemical transport, assimilation, and climate models [*Intergovernmental Panel on Climate Change (IPCC)*, 2001].

[7] In the last decade, research of the global aerosol system was invigorated by an array of new measurements in several temporal and spatial scales, the fruits of which we start to see in this special issue, and the development of new models:

[8] 1. Global, daily, and multidaily observations of aerosol, clouds, and their properties derived from satellite measurements of the scattered solar light, utilizing its spectral, angular, and polarization properties.

[9] 2. Satellite measurements of the vertical distribution of aerosol, clouds, and precipitation using lidars and radars and thermal infrared and microwave radiometers.

[10] 3. Surface networks that continuously measure in situ and remotely (both passive radiometers and active sensors) the aerosol physical and chemical properties, distribution of solar radiation, clouds, and precipitation.

[11] 4. Intensive field campaigns that concentrate the measurement power in a given region for a limited period of time and help understand the aerosol sources and aerosol and cloud processes.

[12] 5. Laboratory analysis of aerosol particles and measurements of aerosol processes.

[13] 6. Development of transport models that start to mimic the aerosol distribution measured from satellite and surface networks.

[14] 7. Assimilation of the aerosol models and measurements into climate models to study the effects of aerosol on

¹Laboratoire d'Optique Atmosphérique, Villeneuve d'Ascq, France.

²NASA Goddard Space Flight Center, Greenbelt, Maryland, USA.

³Center for Climate System Research, University of Tokyo, Tokyo, Japan.

⁴Center for Clouds, Chemistry and Climate, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California, USA.

global circulation and climate and the feedback from climate change on the aerosol.

[15] These new measurements and models are now starting to develop new understanding. The papers in this special issue summarize some of the progress in the following fields.

[16] 1. Remote sensing of aerosol and validation using the new satellite instruments: POLDER [Herman *et al.*, 2005], Moderate Resolution Imaging Spectroradiometer (MODIS) [Ichoku *et al.*, 2005], Multiangle Imaging Spectroradiometer (MISR) [Kahn *et al.*, 2005], Global Ozone Monitoring Experiment (GOME) [Kusmierczyk-Michulec and de Leeuw, 2005], Sea-viewing Wide Field-of-view Sensor (SeaWiFS) [Wang *et al.*, 2005], combining several instruments like MISR and MODIS [Abdou *et al.*, 2005] or including satellites with a long record such as advanced very high resolution radiometer (AVHRR) and Total Ozone Mapping Spectrometer (TOMS) [Jeong and Li, 2005], or MODIS and AVHRR [Jeong *et al.*, 2005].

[17] 2. Preparation for remote sensing with the to be launched satellites that will be part of the A-train POLDER+ CALIPSO [Waquet *et al.*, 2005; C. Catrall *et al.*, Variability of aerosol lidar, backscatter and extinction ratios of key aerosol types derived from selected AERONET locations, submitted to *Journal of Geophysical Research*, 2005].

[18] 3. Use of the new satellite and ground network data with models to infer regional and global aerosol distribution and impacts dust transport and deposition [Kaufman *et al.*, 2005; Liu *et al.*, 2005], regional and global aerosol burden [Massie *et al.*, 2004; Omar *et al.*, 2005; Liu *et al.*, 2005; Reddy *et al.*, 2005], aerosol global absorption [Schuster *et al.*, 2005; Torres *et al.*, 2005], derivation in conjunction with CO measurements [Edwards *et al.*, 2004], and interaction between aerosol and clouds [Krüger *et al.*, 2004; M. Jin *et al.*, Urban aerosols and their variations with clouds and rainfall: A case study for New York and Houston, submitted to *Journal of Geophysical Research*, 2005; V. N. Kapustina *et al.*, On the determination of CCN from satellite: Challenges and possibilities, submitted to *Journal of Geophysical Research*, 2005].

[19] 4. Inferring the aerosol radiative forcing of climate [Vant-Hull *et al.*, 2005; Kim *et al.*, 2005; Zhang *et al.*, 2005a, 2005b].

[20] We can anticipate that in the next several years the data from the present and newly launched satellites will come into full fruition. Plans are underway for several new spaceborne missions for advanced aerosol, clouds, and precipitation measurements, increasing the spectral, angular, and polarization resolution, and introducing new active radar and lidar measurements. Data from these mission, if successfully executed together with developments in atmospheric modeling, are expected to advance our understanding of the interaction of aerosols, clouds, and precipitation and their effect on the hydrological cycle and climate.

References

- Abdou, W. A., D. J. Diner, J. V. Martonchik, C. J. Bruegge, R. A. Kahn, B. J. Gaitley, K. A. Crean, L. A. Remer, and B. Holben (2005), Comparison of coincident Multiangle Imaging Spectroradiometer (MISR) and Moderate Resolution Imaging Spectroradiometer (MODIS) aerosol optical depths over land and ocean scenes containing Aerosol Robotic Network (AERONET) sites, *J. Geophys. Res.*, doi:10.1029/2004JD004693, in press.
- Edwards, D. P., *et al.* (2004), Observations of carbon monoxide and aerosols from the Terra satellite: Northern Hemisphere variability, *J. Geophys. Res.*, 109, D24202, doi:10.1029/2004JD004727.
- Herman, M., J.-L. Deuzé, A. Marchand, B. Roger, and P. Lallart (2005), Aerosol remote sensing from POLDER/ADEOS over the ocean: Improved retrieval using a nonspherical particle model, *J. Geophys. Res.*, 110, D10S02, doi:10.1029/2004JD004798.
- Ichoku, C., L. A. Remer, and T. F. Eck (2005), Quantitative evaluation and intercomparison of morning and afternoon MODIS aerosol measurements from Terra and Aqua satellites, *J. Geophys. Res.*, 110, D10S03, doi:10.1029/2004JD004987.
- Intergovernmental Panel on Climate Change (IPCC) (2001), *Climate Change 2001: The Scientific Basis: Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by J. T. Houghton *et al.*, Cambridge Univ. Press, New York.
- Jeong, M.-J., and Z. Li (2005), Quality, compatibility, and synergy analyses of global aerosol products of global aerosol products derived from the advanced very high resolution radiometer and Total Ozone Mapping Spectrometer, *J. Geophys. Res.*, 110, D10S08, doi:10.1029/2004JD004647.
- Jeong, M.-J., Z. Li, D. A. Chu, and S.-C. Tsay (2005), Quality and compatibility analyses of global aerosol products derived from the advanced very high resolution radiometer and Moderate Resolution Imaging Spectroradiometer, *J. Geophys. Res.*, 110, D10S09, doi:10.1029/2004JD004648.
- Kahn, R. A., B. J. Gaitley, J. V. Martonchik, D. J. Diner, K. A. Crean, and B. Holben (2005), Multiangle Imaging Spectroradiometer (MISR) global aerosol optical depth validation based on 2 years of coincident Aerosol Robotic Network (AERONET) observations, *J. Geophys. Res.*, 110, D10S04, doi:10.1029/2004JD004706.
- Kaufman, Y. J., D. Tanré, and O. Boucher (2002), A satellite view of aerosols in the climate system, *Nature*, 419, 215–223.
- Kaufman, Y. J., I. Koren, L. A. Remer, D. Tanré, P. Ginoux, and S. Fan (2005), Dust transport and deposition observed from the Terra-Moderate Resolution Imaging Spectroradiometer (MODIS) spacecraft over the Atlantic Ocean, *J. Geophys. Res.*, 110, D10S12, doi:10.1029/2003JD004436.
- Kim, D.-H., B.-J. Sohn, T. Nakajima, and T. Takamura (2005), Aerosol radiative forcing over east Asia determined from ground-based solar radiation measurements, *J. Geophys. Res.*, 110, D10S22, doi:10.1029/2004JD004678.
- Krüger, O., R. Marks, and H. Graßl (2004), Influence of pollution on cloud reflectance, *J. Geophys. Res.*, 109, D24210, doi:10.1029/2004JD004625.
- Kusmierczyk-Michulec, J., and G. de Leeuw (2005), Aerosol optical thickness retrieval over land and water using GOME data, *J. Geophys. Res.*, 110, D10S05, doi:10.1029/2004JD004780.
- Liu, H., R. T. Pinker, and B. N. Holben (2005), A global view of aerosols from merged transport models, satellite, and ground observations, *J. Geophys. Res.*, 110, D10S15, doi:10.1029/2004JD004695.
- Massie, S. T., O. Torres, and S. J. Smith (2004), Total Ozone Mapping Spectrometer (TOMS) observations of increases in Asian aerosol in winter from 1979 to 2000, *J. Geophys. Res.*, 109, D18211, doi:10.1029/2004JD004620.
- Omar, A. H., J.-G. Won, D. M. Winker, S.-C. Yoon, O. Dubovik, and M. P. McCormick (2005), Development of global aerosol models using cluster analysis of AERONET measurements, *J. Geophys. Res.*, 110, D10S14, doi:10.1029/2004JD004874.
- Ramanathan, V., P. J. Crutzen, J. T. Kiehl, and D. Rosenfeld (2001), Aerosols, climate, and the hydrological cycle, *Science*, 294, 2119–2124.
- Reddy, M. S., O. Boucher, N. Bellouin, M. Schulz, Y. Balkanski, J.-L. Dufresne, and M. Pham (2005), Estimates of global multicomponent aerosol optical depth and direct radiative perturbation in the LMDZT general circulation model, *J. Geophys. Res.*, doi:10.1029/2004JD004757, in press.
- Schuster, G. L., O. Dubovik, B. N. Holben, and E. E. Clothiaux (2005), Inferring black carbon content and specific absorption from Aerosol Robotic Network (AERONET) aerosol retrievals, *J. Geophys. Res.*, 110, D10S17, doi:10.1029/2004JD004548.
- Torres, O., P. K. Bhartia, A. Sinyuk, E. J. Welton, and B. Holben (2005), Total Ozone Mapping Spectrometer measurements of aerosol absorption from space: Comparison to SAFARI 2000 ground-based observations, *J. Geophys. Res.*, 110, D10S18, doi:10.1029/2004JD004611.
- Vant-Hull, B., Z. Li, B. F. Taubman, R. Levy, L. Marufu, F.-L. Chang, B. G. Doddridge, and R. R. Dickerson (2005), Smoke over haze: Comparative analysis of satellite, surface radiometer, and airborne in situ measurements of aerosol optical properties and radiative forcing over the eastern United States, *J. Geophys. Res.*, 110, D10S21, doi:10.1029/2004JD004518.

- Waquet, F., J.-F. Léon, P. Goloub, J. Pelon, D. Tanré, and J.-L. Deuzé (2005), Maritime and dust aerosol retrieval from polarized and multi-spectral active and passive sensors, *J. Geophys. Res.*, *110*, D10S10, doi:10.1029/2004JD004839.
- Wang, M., K. D. Knobelspiesse, and C. R. McClain (2005), Study of the Sea-Viewing Wide Field-of-View Sensor (SeaWiFS) aerosol optical property data over ocean in combination with the ocean color products, *J. Geophys. Res.*, *110*, D10S06, doi:10.1029/2004JD004950.
- Zhang, J., S. A. Christopher, L. A. Remer, and Y. J. Kaufman (2005a), Shortwave aerosol radiative forcing over cloud-free oceans from Terra: 1. Angular models for aerosols, *J. Geophys. Res.*, *110*, D10S23, doi:10.1029/2004JD005008.
- Zhang, J., S. A. Christopher, L. A. Remer, and Y. J. Kaufman (2005b), Shortwave aerosol radiative forcing over cloud-free oceans from Terra: 2. Seasonal and global distributions, *J. Geophys. Res.*, *110*, D10S24, doi:10.1029/2004JD005009.